Connecticut Department of Public Health HIV Surveillance Program Recently Diagnosed HIV cases by Risk, Sex, Race, and Age Group, Bridgeport, 2014 - 2018

				Mode of Transmission											
	Total		MSM		PWID		MSM and IDU		Heterosexual contact		Presumed Heterosexual contact		Unknown		
	N	% of total	N	% of row total	N	% of row total	N	% of row total	N	% of row total	N	% of row total	N	% of row total	
Total	161	100.0	67	41.6	9	5.6	1	0.6	57	35.4	8	5.0	19	11.8	
Sex															
Male	103	64.0	67	65.0	4	3.9	1	1.0	17	16.5	0	0	14	13.6	
Female	58	36.0	0	0	5	8.6	0	0	40	69.0	8	13.8	5	8.6	
Race/ethncity															
Black/African American	95	59.0	32	33.7	5	5.3	1	1.1	43	45.3	6	6.3	8	8.4	
Hispanic/Latino	43	26.7	22	51.2	3	7.0	0	0	10	23.3	1	2.3	7	16.3	
White	18	11.2	11	61.1	1	5.6	0	0	2	11.1	1	5.6	3	16.7	
Other races	5	3.1	2	40.0	0	0	0	0	2	40.0	0	0	1	20.0	
Age at diagnosis															
<20	5	3.1	2	40.0	1	20.0	0	0	0	0	1	20.0	1	20.0	
20-29	54	33.5	37	68.5	0	0	1	1.9	11	20.4	1	1.9	4	7.4	
30-39	35	21.7	15	42.9	3	8.6	0	0	13	37.1	1	2.9	3	8.6	
40-49	32	19.9	6	18.8	1	3.1	0	0	18	56.3	3	9.4	4	12.5	
50+	35	21.7	7	20.0	4	11.4	0	0	15	42.9	2	5.7	7	20.0	

This table presents HIV Infection cases using the year the person was first diagnosed. The term HIV Infection is used to refer to three categories of diagnoses collectively:

2 - a diagnosis of HIV infection and a later diagnosis of AIDS

3 - concurrent diagnoses of HIV and AIDS.

Source: HIV Surveillance Registry for cases reported through December 2019

^{1 -} persons with a diagnosis of HIV infection (not AIDS)