

PCMH+ Measure List
January 10, 2017

Measure Name	Description	Steward	NQF #	PCMH+ Measure Type
Adolescent well-care visits	This measure is used to assess the percentage of enrolled members 12 through 21 years of age who had at least one comprehensive well-care visit with a primary care practitioner (PCP) or an obstetrics and gynecology (OB/GYN) practitioner during the measurement year.	NCQA		Scoring
Annual fluoride treatment ages 0<4	Annual fluoride treatment ages 0<4 (in a pediatric or dental setting)	DSS		Reporting Only
Annual monitoring for persistent medications (roll-up)	The percentage of patients 18 years of age and older who received a least 180 treatment days of ambulatory medication therapy for a select therapeutic agent during the measurement year and at least one therapeutic monitoring event for the therapeutic agent in the measurement year. - Angiotensin converting enzyme (ACE) inhibitors or angiotensin receptor blockers (ARB) - Digoxin - Diuretics <u>- Total Rate will be measured</u>	NCQA	2371	Reporting Only
Appropriate treatment for children with upper respiratory infection	Percentage of children 3 months to 18 years of age with a diagnosis of URI who were not dispensed an antibiotic medication. A higher rate indicates appropriate care (i.e. the proportion for whom antibiotics were not prescribed)	NCQA	0069	Reporting Only
Asthma Medication Ratio	The percentage of members 5-64 years of age with persistent asthma and had a ratio of controller medications to total medications of 0.50 or greater during the measurement year.	NCQA	1800	Reporting Only
Avoidance of antibiotic treatment in adults with acute bronchitis	The percentage of adults 18–64 years of age with a diagnosis of acute bronchitis who were not dispensed an antibiotic prescription. A higher rate indicates appropriate care (i.e. the proportion for whom antibiotics were not prescribed)	NCQA	0058	Scoring
Behavioral Health Screening 1-17	The percentage of children ages 1-17, who were screened for developmental or behavioral problems using a validated survey instrument, approved by the AAP.	DSS		Challenge
Breast cancer screening	The percentage of women 50-74 years of age who had a mammogram to screen for breast cancer in a two year period.	NCQA	2372	Reporting Only

PCMH+ Measure List
January 10, 2017

Measure Name	Description	Steward	NQF #	PCMH+ Measure Type
Cervical cancer screening	Percentage of women 21–64 years of age who were screened for cervical cancer using either of the following criteria: - Women age 21–64 who had cervical cytology performed every 3 years. - Women age 30–64 who had cervical cytology/human papillomavirus (HPV) co-testing performed every 5 years.	NCQA	0032	Reporting Only
Chlamydia screening in women	The percentage of women 16–24 years of age who were identified as sexually active and who had at least one test for chlamydia during the measurement year.	NCQA	0033	Reporting Only
Developmental screening in the first three years of life. Three age breakouts (ages 1, 2, and 3)	The percentage of children screened for risk of developmental, behavioral and social delays using a standardized screening tool in the first three years of life. This is a measure of screening in the first three years of life that includes three, age-specific indicators assessing whether children are screened by 12 months of age, by 24 months of age and by 36 months of age. (The total of the three ages will be scored)	OHSU	1448	Scoring
Diabetes eye exam	The percentage of patients 18-75 years of age with diabetes (type 1 and type 2) who had at least one eye exam (retinal) performed in a two year period.	NCQA	0055	Reporting Only
Diabetes HbA1c Screening	Adults age 18-75 with a diagnosis of Type I or Type II diabetes who received at least one HbA1c screening during the measurement year.	NCQA	0057	Scoring
Diabetes: medical attention for nephropathy	The percentage of patients 18-75 years of age with diabetes (type 1 and type 2) who received a nephropathy screening test or had evidence of nephropathy during the measurement year.	NCQA	0062	Reporting Only
ED Usage	Emergency department usage (Excludes mental health and chemical dependency services).	NCQA		Scoring
Follow-up care for children prescribed ADHD medication	The percentage of children ages 6-12 as of the Index Prescription Start Date (IPSD) newly prescribed attention-deficit/hyperactivity disorder (ADHD) medication who had at least three follow-up care visits within a 10-month period, one of which was within 30 days of when the first ADHD medication was dispensed. Two rates are reported: <ul style="list-style-type: none"> • Initiation Phase and Continuation Phase 	NCQA	0108	Reporting Only

PCMH+ Measure List
January 10, 2017

Measure Name	Description	Steward	NQF #	PCMH+ Measure Type
Human Papillomavirus Vaccine for Adolescents (HPV)*	The percentage of adolescents 13 years of age who had three doses of the HPV vaccine by their 13th birthday.*	NCQA	1959	Reporting Only
Medication management for people with asthma	Medication Management for people with asthma age 5-64 (age 5-18 breakout can be used for pediatric practices). Percent of patients with <i>persistent</i> asthma who were prescribed and remained on asthma "controller medication" for at least 75% of their treatment period.	NCQA	1799	Scoring
Metabolic Monitoring for Children and Adolescents on Antipsychotics	Percentage of children and adolescents 1-17 years of age who had two or more antipsychotic prescriptions and had metabolic testing	NCQA		Challenge
Oral evaluation, dental services	Percentage of enrolled children under age 21 who received a comprehensive or periodic oral evaluation within the reporting year	American Dental Association	2517	Reporting Only
PCMH CAHPS	Consumer Assessment of Healthcare Providers and Systems ® CAHPS - PCMH version.		N/A	Scoring
Post-Hospital Admission Follow-up	Percentage of adults age 21-75 with an inpatient "medical" or psych admission with a claim for post-admission follow-up with a physician, PA, or APRN within seven days of the inpatient discharge. Medical admissions are defined as all admissions that are not maternity or surgery related.	DSS		Challenge

PCMH+ Measure List
January 10, 2017

Measure Name	Description	Steward	NQF #	PCMH+ Measure Type
Prenatal care & Postpartum care	<p>The percentage of deliveries of live births between November 6 of the year prior to the measurement year and November 5 of the measurement year. For these women, the measure assesses the following facets of prenatal and postpartum care.</p> <ul style="list-style-type: none"> • Rate 1: Timeliness of Prenatal Care. The percentage of deliveries that received a prenatal care visit as a patient of the organization in the first trimester or within 42 days of enrollment in the organization. • Rate 2: Postpartum Care. The percentage of deliveries that had a postpartum visit on or between 21 and 56 days after delivery. 	NCQA	1517	Scoring
Readmission	Readmissions Within 30 Days (PH and BH)	MMDN		Challenge
Use of imaging studies for low back pain	The percentage of members with a primary diagnosis of low back pain who did not have an imaging study (plain x-ray, MRI, CT scan) within 28 days of the diagnosis.	NCQA	0052	Reporting Only
Well-child visits in the first 15 months of life	<p>Percentage of patients who turned 15 months old during the measurement year and who had the following number of well-child visits with a PCP during their first 15 months of life.</p> <ul style="list-style-type: none"> •Six or more well-child visits 	NCQA	1392	Scoring
Well-child visits in the third, fourth, fifth and sixth years of life	Percentage of patients 3–6 years of age who received one or more well-child visits with a PCP during the measurement year.	NCQA	1516	Reporting Only

* Males have been added to the 2017 HEDIS Measure